RADIOACTIVE ROBINS

photography: **NORMAN DEANS VAN SWELM**

2009

1 Waders - Kentish Plover

Strandplevier

[1 Kentish Plover] [1 Eurasian Whimbrel] [1 Eurasian Curlew] [1 Common Redshank] [1 Ruddy Turnstone] [1 Sanderling] [1 Dunlin] [1 Ruff] [1 Common Snipe] [1 Jack Snipe] [1 Eurasian Woodcock] [1 Black-tailed Godwit] [1 Bar-tailed Godwit] [1 Eurasian Whimbrel] [1 Little Stint] [1 Pectoral Sandpiper] [1 Common Redshank] [1 Wood Sandpiper] [1 Green Sandpiper] [1 Marsh Sandpiper] [1 Common Greenshank] [1 Spotted Redshank] [1 Ruddy Turnstone] [1 Red Knot] [1 Sanderling] [1 Dunlin] [1 Ruff] [1 Stilt Sandpiper] [1 Buff-breasted Sandpiper] [1 Pectoral Sandpiper] [1 Curlew Sandpiper] [1 Jack Snipe] [1 Eurasian Woodcock] [1 Black-tailed Godwit] [1 Bar-tailed Godwit] [1 Eurasian Whimbrel] [1 Eurasian Curlew] [1 Little Stint] [1 Temminck's Stint] [1 Pectoral Sandpiper

2 Waders - Hudsonian Whimbrel

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

Hudsonian Whimbrel *Numenius (p)h.hudsonicus* Engelmoer & Roselaar (1998) recognize two subspecies of Nearctic Whimbrel *Numenius (p)hudsonicus* i.e. nominate *hudsonicus* as well as *rufiventris*. *Rufiventris*, which breeds from W. Alaska east to the NW territories, has the typical brownish appearance as in the bird in the first picture.
Nominate *hudsonicus* breeds along the south and west coast of the Hudson Bay.
*Engelmoer, Meinte and Cees S.Roselaar. 1998. Geographical Variation in Waders. Kluwer Academic Publishers. Dordrecht/Boston/London: ISBN 0-7923-5020-0



American Whimbrel Numenius hudsonicus dec.1976 Galapagos

Hudsonian Whimbrel Numenius h.hudsonicus Dec.1976 Galapagos

NAVIGATE

08 feb 2009

3 Waders - Hudsonian Dunlin

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

This Hudsonian Dunlin Calidris (alpina) hudsonia discovered on 24th May 2001 by Charles Deans van Swelm along the south bank of the Wester Scheldt near Terneuzen in The Netherlands is the first record for the Western Palearctic. Recently another Hudsonian Dunlin was observed in Europe, this time a 1st winter bird on the Isles of Scilly, Britain on 29th October 2007 (Birding World 20 (11): 464-466.

Hudsonian Dunlin C.a.hudsonia 24052001 Wester Scheldt (W-Schelde), Terneuzen, The Netherlands

NAVIGATE

08 feb 2009

4 Waders - Interesting Wader

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

Interesting Wader On the 21st February 2008 this wader (1, 2, 3) was briefly seen at a high tide roost on the beach of Oostvoorne, The Netherlands. It's jizz suggests White-Rumped Sandpiper *Calidris fuscicollis*. The bird seems in active wing-moult which would fit White-rumped Sandpiper well as it moults during November-March. All descriptions known to us however mention grey-brown or pale brownish streaks on breast and/or flanks in both adult winter and 1st winter plumage which this birds clearly does not show. We would like to know if anyone is familiar with this plumage.

wader spec.21022008 3 Oostvoorne, The Netherlands

wader spec.21022008 1 Oostvoorne, The Netherlands

wader spec.21022008 2 Oostvoorne, The Netherlands

NAVIGATE

08 feb 2009

5 Waders - Knot *Calidris canutus*

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

Knot *Calidris canutus* (see pictures below)

Standing knowledge is that Knots wintering in NW Europe belong to the Greenland race *islandica*. Adult birds return after breeding to NW Europe from the end of July onwards and undergo a complete moult in the months that follow. Juveniles arrive from the latter part of August onwards. The Waddensea in The Netherlands and Germany is of crucial importance for this population. Greenland Knots especcially have suffered greatly from mechanical Cockle fisheries which is now largely forbidden. Greenland Knots leave Holland in the 2nd half of April when still in winter plumage. Nominate *canutus* , the Siberian Knot, breeds in Russia on the Taimyr peninsula and winters in West Africa, in particular on the Banc d'Arguin in Mauritania. These birds use the Waddensea as a staging area on the way to or back from Taimyr to 'refuel' used up fat supplies during May and August. When they arrive in spring they are already in breeding plumage.

Knots *Calidris canutus* in 1st summer plumage (see Waders - Knots page 5b)

The Knots wintering in the Rhine-Meuse delta in the Netherlands belong to the Greenland race *islandica*. By the end of April they look fat & are about to leave for the breeding grounds. safe for a small number of 2nd calendar year birds, these young birds begin moulting their wing 1st winter plumage into their 2nd winter plumage as can be seen on the pictures.

Knot *C.c.islandica* 21122007 Oostvoorne, The Netherlands



01Knots Calidris c.islandica 27022008 1 Oostvoorne, The Netherlands



02Knots Calidris c.islandica 27022008 2 Oostvoorne, The Netherlands



03Knots Calidris c.islandica 27022008 3 Oostvoorne, The Netherlands



04Knots Calidris c.islandica 27022008 4 Oostvoorne, The Netherlands

05Knots Calidris c.islandica 27022008 5 Oostvoorne, The Netherlands

06Knots Calidris c.islandica 27022008 6 Oostvoorne, The Netherlands



07Knots Calidris c.islandica 27022008 7 Oostvoorne, The Netherlands

08Knots Calidris c.islandica 27022008 8 Oostvoorne, The Netherlands

09Knots Calidris c.islandica in flight 27022008 Oostvoorne, The Netherlands

to 5b> Knots *Calidris canutus* in 1st summer plumage (see Waders - Knots page 5b)

The Knots wintering in the Rhine-Meuse delta in The Netherlands belong to the Greenland race *islandica*. By the end of April they look fat & are about to leave for the breeding grounds. safe for a small number of 2nd calendar year birds, these young birds begin moulting their wing 1st winter plumage into their 2nd winter plumage as can be seen on the pictures.

NAVIGATE

08 feb 2009

6 Waders - Baltic Dunlin

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

The Baltic Dunlin *Calidris alpina schinzii* is the rarest of the Dunlin races in Europe.
It's world population is estimated to be no more than 6,000 individuals. It has occassionally bred in The Netherlands. It is this Dunlin that lives after it's nickname 'Plover's page". In the Dutch delta area it can be found in groups of Ringed Plover *Charadrius hiaticula* during spring migration (February-March), where they occur on the higher, more vegetated parts of the tidal areas. They keep well apart from the more numerous Northern Dunlin *Calidris alpina alpina* which occurs in the lower muddier parts. The Baltic Dunlin is the first to complete it's breeding plumage (March).

Calidris alpina schinzii 19022008 Oostvoorne, The Netherlands

NAVIGATE

08 feb 2009

7 Waders - The Sanderling Calidris alba

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

1.Sanderlings Calidris alba braving the storm 20032008 Brouwersdam, The Netherlands

2.Sanderlings braving the storm, head formation 20032008

3.Sanderlings braving the storm, life in the middle 20032008

4.Sanderlings storm formation, tail section 20032008

5.Sanderlings braving the storm, changing formation 20032008

6.Sanderlings braving the storm on your own 20032008

7.Sanderlings braving the storm & enduring the misery 20032008

NAVIGATE

08 feb 2009

8 Little Stint *Calidris minuta*

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

Little Stint *Calidris minuta*

As the main route during spring migration from Africa, where this small wader stays during winter, to the northern tundra's,where it breeds, leads over the middle of the Mediterranean and farther east only small numbers reach the North Sea coast in May. They arrive in breeding plumage which they obtained during a partial moult in Africa.

Little Stint Calidris minuta 02052008a Slikken van Flakkee, The Netherlands

Little Stint Calidris minuta 02052008b Slikken van Flakkee, The Netherlands

NAVIGATE

08 feb 2009

9. Eskimo Curlew *Numenius borealis*

Species Status List | Species Status List Search | Species Status List Menu | Species Status List Table Front | Species Status List Table Back | Species Status List Table Back 2 | Species Status List Table Back 3 | Species Status List Table Back 4 | Species Status List Table Back 5 | Species Status List Table Back 6 | Species Status List Table Back

10 Dunlins *Calidris alpina* in juvenile plumage

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint Calidris minuta> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin Calidris alpina in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons>

dunlin calidris alpina/Baltic Dunlin C.a.schinzii juv 14072008 Rotterdam, The Netherlands

dunlin calidris alpina/Baltic Dunlin C.a.schinzii juv back 14072008 Rotterdam, The Netherlands

dunlin calidris alpina/Baltic Dunlin C.a.schinzii juv flight 14072008 Rotterdam, The Netherlands

dunlin calidris alpina/Icelandic Dunlin C.a.islandica juv. 22072008 1 Brouwersdam, The Netherlands

dunlin calidris alpina/Icelandic Dunlin C.a.islandica juv. 22072008 2 Brouwersdam, The Netherlands

dunlin calidris alpina/Icelandic Dunlin C.a.islandica juv. 22072008 3 Brouwersdam, The Netherlands

dunlin calidris alpina/Icelandic Dunlin C.a.islandica juv. 22072008 4 Brouwersdam, The Netherlands

dunlin calidris alpina/Icelandic Dunlin C.a.islandica juv. 22072008 Brouwersdam, The Netherlands

dunlin calidris alpina/Dunlin Calidris a.alpina juv. 21082008 Rotterdam, The Netherlands

dunlin calidris alpina/Dunlin Calidris a.alpina juv 21082008 1 Rotterdam, The Netherlands

dunlin calidris alpina/Dunlin C.a.alpina juv 01092008 Roompotsluizen, The Netherlands

11 Whimbrel Numenius phaeopus

Taxonomy & Species: CHRS (Sibley/Ahlquist taxonomy) • Species: Whimbrel Numenius phaeopus • Subspecies: Nominate phaeopus (Eurasian Whimbrel) • Alternate Common Name: Eurasian Whimbrel • Alternate Scientific Name: Numenius phaeopus phaeopus

Whimbrel Numenius phaeopus 19072003 Rotterdam, The Netherlands

Whimbrel Numenius phaeopus 19072003 Rotterdam, The Netherlands

Whimbrel Numenius phaeopus 22072008 Oudeloxy, The Netherlands

Whimbrel Numenius phaeopus 22072008 Oudeloxy, The Netherlands

Whimbrel Numenius phaeopus 22072008 Brouwersdam, The Netherlands

Whimbrel Numenius phaeopus 22072008 Oudeloxy, The Netherlands

Whimbrel Numenius phaeopus 22072008 Oudeloxy, The Netherlands

Whimbrel Numenius phaeopus 22072008 Brouwersdam, The Netherlands

Whimbrel Numenius phaeopus ad 07082008 Brouwersdam, The Netherlands

Whimbrel Numenius phaeopus 07082008 Rotterdam, The Netherlands

Whimbrel Numenius phaeopus ad 07082008 Rotterdam, The Netherlands

Whimbrel Numenius phaeopus 07082008 Rotterdam, The Netherlands

Whimbrel Numenius phaeopus 14082007 Oostvoorne, The Netherlands

Whimbrel Numenius phaeopus 14082007 Oostvoorne, The Netherlands

a/b Whimbrel Numenius phaeopus juvenile 16062008 Balkema v Adrian Dammenoud Hill

Whimbrel Numenius phaeopus ad 20042008 Rotterdam, The Netherlands

Whimbrel Numenius p.variegatus 01102008 Hong Kong v Neil Fifer

Whimbrel Numenius phaeopus juvenile 04102004 Rotterdam, The Netherlands

Antiquiet

12 Waders - Golden Plover

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

Golden Plover Pluvialis apricaria 1W 25092008 Oostvoorne a, The Netherlands

Golden Plover Pluvialis apricaria 1W 25092008 Oostvoorne b, The Netherlands

NAVIGATE

08 feb 2009

13 Waders - Cannibalism in the cold

1 Kentish Plover> / 2 Hudsonian Whimbrel> / 3 Hudsonian Dunlin> / 4 Interesting Wader> / 5 Knots> / 6 Baltic Dunlin> / 7 The Sanderling Calidris alba> / 8 Little Stint *Calidris minuta*> / 9 Eskimo Curlew Numenius borealis> / 10 Dunlin *Calidris alpina* in juvenile plumage> / 11 Whimbrel Numenius phaeopus> / 12 Golden Plover> / 13 Cannibalism in the cold> / 14 A plumage for all seasons> /

During the very cold spell this winter which started on Boxing Day 2008, waders in the Dutch Rhine-Meuse delta began to perish after ten days of severe frost with the lowest recorded temperature of -22C in the SE of the country. As expected Oystercatchers *Haematopus ostralegus* were the first waders to die. On 11th January a very hungry Oystercatcher was found while eating a frozen member of it's own species.

Starved Oystercatcher Haematopus ostralegus eating from the same species corpse

11012009 262 Stellendam, The Netherlands

NAVIGATE

08 feb 2009

14 Waders-A Plumage for all seasons

1 Knight Plover(1) **2** Pacific Golden Plover(2) **3** Grey Plover(3) **4** Northern Sandy(4) **5** Greenshank(5) **6** Common Redshank(6) **7** Spotted Redshank(7) **8** Common Sandpiper(8) **9** Green Sandpiper(9) **10** Wood Sandpiper(10) **11** Terek Sandpiper(11) **12** Marsh Sandpiper(12) **13** Common Snipe(13) **14** Pintail Snipe(14) **15** Jack Snipe(15) **16** Black-tailed Godwit(16) **17** Bar-tailed Godwit(17) **18** Eurasian Woodcock(18) **19** Little Ringed Plover(19) **20** Kentish Plover(20) **21** Common Ringed Plover(21) **22** Eurasian Dotterel(22) **23** Eurasian Oystercatcher(23) **24** Eurasian Curlew(24) **25** Whimbrel(25) **26** Black-winged Stilt(26) **27** Pied Avocet(27) **28** Red-necked Phalarope(28) **29** Grey Phalarope(29) **30** Ruff(30) **31** Little Stint(31) **32** Temminck's Stint(32) **33** Long-toed Stint(33) **34** Sharp-tailed Sandpiper(34) **35** Dunlin(35) **36** Curlew Sandpiper(36) **37** Purple Sandpiper(37) **38** Dunlin(38) **39** Red Knot(39) **40** Great Knot(40) **41** Sanderling(41) **42** Ruddy Turnstone(42) **43** Black Turnstone(43) **44** Eurasian Oyster Catcher(44) **45** Pied Oyster Catcher(45) **46** Black Oyster Catcher(46) **47** Crab Plover(47) **48** Eurasian Thick-knee(48) **49** Cream-coloured Courser(49) **50** Collared Pratincole(50) **51** Oriental Pratincole(51) **52** Wheatear(52) **53** Northern Wheatear(53) **54** Isabelline Wheatear(54) **55** Desert Wheatear(55) **56** Pied Wheatear(56) **57** Black Wheatear(57) **58** Stone Chat(58) **59** Pied Bush Chat(59) **60** Grey Bush Chat(60) **61** Rock Thrush(61) **62** Blue Rock Thrush(62) **63** Blue Chat(63) **64** Bluethroat(64) **65** Red-breasted Flycatcher(65) **66** Pied Flycatcher(66) **67** Red-breasted Flycatcher(67) **68** Red-breasted Flycatcher(68) **69** Red-breasted Flycatcher(69) **70** Red-breasted Flycatcher(70) **71** Red-breasted Flycatcher(71) **72** Red-breasted Flycatcher(72) **73** Red-breasted Flycatcher(73) **74** Red-breasted Flycatcher(74) **75** Red-breasted Flycatcher(75) **76** Red-breasted Flycatcher(76) **77** Red-breasted Flycatcher(77) **78** Red-breasted Flycatcher(78) **79** Red-breasted

This website is dedicated to

Charles Darwin (1809-1882)

The greatest naturalist there ever was!>

http://home.planet.nl/~swelm001/

Charles Darwin

LAST UPDATED 08 feb 2009

PL

RADIOACTIVEROBINS.COM

http://www.radioactiverobins.com

p p

RADIOACTIVE ROBINS

(.....and other birds !)

Foundation Voorne Bird Observatory - The Netherlands

NORMAN DEANS VAN SWELM > ON BIRDS

MAIL

RADIOACTIVE ROBINS>	CHERNOBYL /// downld .pdf about Radioactive Robins (140kb)	
ID-PROBLEMS>	QUESTIONS	
Colour mutation>	Colour mutation	
Abnormal plumages>	Rare case of gynandromorphy	
Hybrids>	Hybrids	
GREBES>	Black-necked Grebe	Horned - or Slavonian Grebe
	Great Crested Grebe	
SEABIRDS>	Arctic Quillemot	Masked Booby
	Blue footed booby	Strange Fulmar
	Cormorants > a>	Wedge-tailed Shearwater
HERONS & EGRETS>	Arabian Reef Heron	Little Egret
	Great Blue Heron	Purple Heron
	Grey Heron	
SWANS & GEESE>	Bewick's Swan	Geese
	Whistling Swan	
DUCKS>	>central-index all waterbirds 0- Canadian Eiders in the North Sea 1-Casaindexducks.htmrca Tadorna ferruginea 2-Teal Anas crecca	3-Gadwall Anas strepera 4-Shelduck 5-Hybrids 6-Oldsquaw Clangula hyemalis hybridisation 7-Shoveler Anas clypeata
BIRDS OF PREY>	Merlin/Falco columbarius	Escaped birds of prey and their origin
	an interesting falcon	how a Peregrine lost it's prey to a Kestrel
	hobby *Falco subbuteo*	
WADERS & SNIPES>	Kentish Plover	Snipe *Gallinago gallinago* & Pin-tailed Snipe *Gallinago stenura*
	Hudsonian Whimbrel	Hudsonian Dunlin
	Knots	Interesting Wader
	Knots in 1st summerplumage	The Sanderling Calidris alba
	Baltic Dunlin	Eskimo Curlew Numenius borealis
	Little Stint Calidris minuta	Dunlins *Calidris alpina* in juvenile plumage
	Golden Plover	Whimbrel Numenius phaeopus
	Waders - Cannibalism in the cold	
SKUA'S> & LARGE GULLS>	Arctic Skua	Heuglin's & Baraba Gull
	Armenian Gull	Iceland Gull
	Atlantic Gull	Kelp Gull/Larus Dominicanus
	Larus atl gibraltar picks olives	Kumlien's Gull
	Caspian- & Marsh gull	Lesser Black-backed Gull - gallery
	Dutch Herring gull	Russian Herring Gull L.a.argentatus
	Glaucous Gull	Taimyr Gull
	Glaucous Winged Gull	Thayer's Gull
	Arctic Gull	Yellow-legged Gull
	Great Black-backed Gull Larus Marinus 1 - 2 - 3 - 4 - 5 - 6	Young herring gull: a never ending puzzle
MEDIUM-SIZED GULLS> & SMALL GULLS>	Black-Headed Gull	Little Gull
	Bonaparte's Gull	Mediterranean Gull & climate change
	Common Gull	Short-billed or Mew Gull
	The Dutch Common Gull	Moongull
	Franklin's Gull Larus pipixcan	Ross' Gull
	Ivory Gull	Russian Common Gull
	Kittiwake	Swallow-tailed Gulls
		Sabine's Gulls
TERNS>	American Black Tern	Garfish pulling
	Arctic Tern	Hybrid Tern
	Common Tern	Little Tern
	Forster's Tern	Sandwich Tern / American
	>>>INDEX TERNS	Sooty Tern on the Island of Eggs
LARKS>	----------	----------
OWLS & NIGHTJARS>	Owls	European Nightjar
		Kingfisher
PIPITS & WAGTAILS>	central-index all wagtails a - White Wagtails b - White Wagtails c - White Wagtails d - Pied Wagtail Motacilla alba yarrellii e - Transbaicalian White Wagtail ee - White wagtail Hybridisation Motacilla alba baicalensis f - Blue-headed Yellow Wagtail Motacilla flava flava - A g - Blue-headed Yellow Wagtail Motacilla flava flava - B h - Blue-headed Yellow Wagtail Motacilla flava flava - C	i - British Yellow Wagtail Motacilla flava flavissima - D j - Grey-headed Yellow Wagtail Motacilla flava thunbergi - E k - Italian OR Ashy-headed Yellow Wagtail Motacilla flava cinereocapilla - F l - Spanish Yellow Wagtail Motacilla flava iberiae - G m - Sykes' Yellow Wagtail *Motacilla flava beema* - H n - Black-headed Wagtail *Motacilla flava feldegg* o - Eastern Yellow Wagtail Motacilla flava tschutschensis - I p - Citrine Wagtails
WHEATEARS>	Red-tailed Wheatear	
ACCENTORS>	Strange Accentor	
OTHER PASSERINES>	British Bullfinches	Redpolls
MAMMELS>	Snow Leopard, near extinct	

All pictures by Norman Deans van Swelm,

unless mentioned otherwise.

Copyright remains with the photographers

MAIL

PL 2007 myspace counter

counter -> started from: may-27-2007

webmaster: rom mager>>>

ro.mar@upcmail.nl

RADIOACTIVE ROBINS.COM

up